

DAMAGE THROUGH DIVISION OF LABOUR: TERMITES AS SOCIAL PESTS

Tian Wu, Vicki Simkovic, Graham J Thompson

Department of Biology, Western University, London, Ontario Canada N6A 3K7

Termites are a relatively small but ancient group of insects that radiated from a cockroach-like ancestor roughly 200 millions years ago (Eggleton, 2011). This evolutionary persistence appears at odds however with the frailty of an individual termite. Termites are soft-bodied, mostly blind, short-lived insects that are prone to desiccation and disease. They make easy prey for ants and other insectivores, and require constant contact and communication with their colony mates for survival. This collective persistence of termite species despite their individual fragility can be explained by two key elements of their biology – namely, a social lifestyle and a propensity to eat wood. The peculiar wood-eating characteristics of termites and their propensity for social living are the very qualities that predispose some species to invade urban, industrial or agricultural habitats as structural pests (Rust & Su, 2012).

As social insects, termites live in kin-based colonies in which individuals are specialized for different roles and depend on each other for survival. The principle division within termite societies is between reproductive and non-reproductive castes, whereby the king and queen are specialized for sex and reproduction, while the soldier and worker castes selflessly labor on behalf of their reproducing relatives. This division in colony labor affords termite colonies an economy of scale that is not available to solitary insects, and it factors into their voraciousness as pests (Wu et al., 2015). For termite societies, the colony collective is therefore more resilient and consuming than the sum of its parts, an element of their social biology that is not lost on pest control companies that seek to systematically control the expansiveness and persistence of invasive colonies (Chouvenc et al., 2011).

In Canada, termite diversity is low but there is one invasion with Toronto as the epicenter. The Eastern subterranean termite *Reticulitermes flavipes* (Kollar) likely invaded Ontario via Toronto in the 1930s, probably from contaminated material shipped from the United States across Lake Ontario (Kirby, 1965). From this point of entry subterranean termites have spread to a vast region that engulfs this Metropolitan city and its provincial surrounds (Raffoul et al., 2011). Moreover, genetic analysis reveals a number of subsequent invasions into Canada, including to Kincardine, Guelph, Leamington and Windsor, among other Ontario municipalities (Scaduto et al., 2012). Finally, unpublished reports suggest new populations have sprung up in other parts of Canada, including Winnipeg (Manitoba) and the Okanagan (British Columbia).

Our lab at Western University is studying the invasive biology of the Eastern subterranean termite. We ask two principle questions. First, how does this species repeatedly invade into, and persist within, Canada, which is just north of its native range.

To answer this question, we use behavioral genetic assays on field-caught termites to test how invasive populations differ from their native counterparts. One principle finding is that invasive populations have very subtle kin recognition (Simkovic et al., 2015), which may explain why colonies invasive to Ontario tend not to assort by kin-colony and instead form massive underground supercolonies that become firmly entrenched and difficult to eradicate. Second, we ask, why upon invasion do termites abandon their discrete colonies in favor of far-reaching supercolonies? To answer this question, we have sequenced the termite transcriptome (all of the protein-encoding regions of the termite genome) from both native and invasive populations, and are using this newly-assembled data set to test a possible genetic explanation (Wu & Thompson, 2015).

We believe it is possible that termite invasions typically involve genetic 'bottlenecks' that strip founding populations of alleles (variants of a gene) that they require to accurately distinguish kin from non-kin. If this reasoning proves correct for termites invasive to Ontario, then we expect low levels of allelic diversity from invasive populations relative to native populations. In the near future, our research could position our lab to isolate the very genes involved in colony formation. We could potentially target them for 'knock down' via genomic technologies emerging from the pest control sector (Scharf, 2015). Given the negative impact that termites continue to bore on wood and wood preservation industries, we see our socio-genetic research on termites at Western University as highly relevant to the Canadian Wood Preservation Association.

Chouvenc, T., Su, N.-Y. & Grace, J.K. (2011) Fifty years of attempted biological control of termites - Analysis of a failure. *Biol. Control.*, 59, 69-82.

Eggleton, P. (2011) An Introduction to Termites: Biology, Taxonomy and Functional Morphology. In Bignell, D.E., Roisin, Y. & Lo, N. (eds), *Biology of Termites: a Modern Synthesis*. Springer, Dordrecht, pp. 1-26.

Kirby, C.S. (1965) The distribution of termites in Ontario after 25 years. *Can. Entomol.*, 97, 310-314.

Raffoul, M., Hecnar, S.J., Prezioso, S., Hecnar, D.R. & Thompson, G.J. (2011) Trap response and genetic structure of Eastern subterranean termites (Isoptera, Rhinotermitidae) in Point Pelee National Park, Ontario, Canada. *Can. Entomol.*, 143, 263-271.

Rust, M.K. & Su, N.Y. (2012) Managing social insects of urban importance. *Annu. Rev. Entomol.*, 57, 355-375.

Scaduto, D.A., Garner, S.R., Leach, E.L. & Thompson, G.J. (2012) Genetic evidence for multiple invasions of the Eastern subterranean termite into Canada. *Environ. Entomol.*, 41, 1680-1686.

Scharf, M.E. (2015) Termites as targets and models for biotechnology. In Berenbaum, M.R. (ed), *Annual Review of Entomology*, Vol 60. pp. 77-102.

Simkovic, V., Thompson, G.J. & McNeil, J.N. (2015) Kin sorting promotes social order in populations of a subterranean termite. *Joint Annual Meeting of the Société d'Entomologie du Québec and the Entomological Society of Canada*. Montreal, Canada.

Wu, T., Simkovic, V. & Thompson, G.J. (2015) Subterranean termites: the evolution of a pest. *PCT Canada*, 3, 34-42.

Wu, T. & Thompson, G.J. (2015) Genomic correlates to kin recognition and invasiveness in a subterranean termite. *Biology and Genomics of Social Insects*. Cold Spring Harbor Laboratories.